


Bougie Through i-gel® Technique for Endotracheal Intubation on Cadavers

Shannon Thompson, MD, MAEd; Jennifer Achay, AS, NRP;
Rachel Ely, DO, MHA, NRP; Stephen Rahm, NRP;
Karl Kmiecik, MD; Eric Elizondo, NRP;
David Wampler, PhD, LP, FAEMS; Craig Navarijo, LP;
Christopher Hewitt, DO

Disclaimer

- The views expressed are those of the presenters and do not reflect the official views or policy of the Department of Defense or its Components.
- The voluntary, fully informed consent of the subjects used in this research was obtained as required by 32 CFR 219 and DODI 3216.02_AFI 40-402.



Background

Hypotheses

The first-pass success rate for endotracheal intubation using a bougie-through i-gel[®] technique will be approximately 70%.


There will be no difference in first-pass success between the uncomplicated or complicated airway.

Methods

- Block randomized crossover study with 2 groups
- Primary outcome was first pass success at intubation
- Complicated airway simulated by c-collar as in a previous study
- Exit survey completed after trials

Results

- 25 paramedics from two fire-based EMS agencies participated.
- The mean years of experience was 4.6 (95% CI 2.3 to 6.8).
- The first-pass success rates for the uncomplicated and complicated airway were 88% (22 of 25) and 76% (19 of 25), respectively.
- Twenty-one (84%) of participants reported 0-5 uses of the i-gel[®] previously.
- There was no difference found in success rate between the uncomplicated and complicated airways ($p=0.45$). There was also no difference in overall success rate between randomization groups ($p=0.467$).



Discussion

Conclusion

- Paramedics using the bougie-through i-gel® technique on cadavers had similar first-pass success rates when compared to previous studies and between uncomplicated and complicated groups.
- The study was not powered sufficiently to compare study first pass success to the population first pass success rate.
- Further research is needed to determine if this technique is applicable to prehospital patients as this was a laboratory study on cadavers.

References

1. Michalek P, Donaldson W, Graham C, Hinds JD. A comparison of the I-gel supraglottic airway as a conduit for tracheal intubation with the intubating laryngeal mask airway: A manikin study. *Resuscitation*. 2010;81(1):74-77. doi:10.1016/j.resuscitation.2009.10.009
2. Kannaujia A, Srivastava U, Saraswat N, Mishra A, Kumar A, Saxena S. A Preliminary Study of I-Gel:A New Supraglottic Airway Device. *Indian J Anaesth*. Published online 2009:5.
3. Melissopoulou T, Stroumpoulis K, Sampanis MA, et al. Comparison of blind intubation through the I-gel and ILMA Fastrach by nurses during cardiopulmonary resuscitation: A manikin study. *Heart Lung*. 2014;43(2):112-116. doi:10.1016/j.hrtlng.2013.12.004
4. Bielski A, Smereka J, Madziala M, Golik D, Szarpak L. Comparison of blind intubation with different supraglottic airway devices by inexperienced physicians in several airway scenarios: a manikin study. *Eur J Pediatr*. 2019;178(6):871-882. doi:10.1007/s00431-019-03345-4
5. Eisenberg MA, Green-Hopkins I, Werner H, Nagler J. Comparison Between Direct and Video-assisted Laryngoscopy for Intubations in a Pediatric Emergency Department. *Acad Emerg Med Off J Soc Acad Emerg Med*. 2016;23(8):870-877. doi:10.1111/acem.13015
6. Halwagi AE, Massicotte N, Lallo A, et al. Tracheal Intubation Through the I-gel™ Supraglottic Airway Versus the LMA Fastrach™: A Randomized Controlled Trial. *Anesth Analg*. 2012;114(1):152–156. doi:10.1213/ANE.0b013e318236f438
7. Theiler LG, Kleine-Brueggeney M, Kaiser D, et al. Crossover Comparison of the Laryngeal Mask Supreme™ and the i-gel™ in Simulated Difficult Airway Scenario in Anesthetized Patients. *Anesthesiol J Am Soc Anesthesiol*. 2009;111(1):55-62. doi:10.1097/ALN.0b013e3181a4c6b9
8. Frerk C, Mitchell VS, McNarry AF, et al. Difficult Airway Society 2015 guidelines for management of unanticipated difficult intubation in adults†. *Br J Anaesth*. 2015;115(6):827-848. doi:10.1093/bja/aev371
9. Campbell J, Michalek P, Deighan M. I-gel supraglottic airway for rescue airway management and as a conduit for tracheal intubation in a patient with acute respiratory failure. *Resuscitation*. 2009;80(8):963. doi:10.1016/j.resuscitation.2009.04.037
10. Gupta L, Chaudhary K, Bhadoria P. I-Gel: A Rescue Intubation Device in Unanticipated Difficult Intubation for Emergency Laparotomy. 2012;2012. doi:10.4236/ojanes.2012.22011

References

11. Gawlowski P, Smereka J, Madziala M, Szarpak L, Frass M, Robak O. Comparison of the Macintosh laryngoscope and blind intubation via the iGEL for Intubation With C-spine immobilization: A Randomized, crossover, manikin trial. *Am J Emerg Med*. 2017;35(3):484-487. doi:10.1016/j.ajem.2016.11.064
12. Bielski A, Rivas E, Ruetzler K, et al. Comparison of blind intubation via supraglottic airway devices versus standard intubation during different airway emergency scenarios in inexperienced hand. *Medicine (Baltimore)*. 2018;97(40). doi:10.1097/MD.00000000000012593
13. Gatward JJ, Cook TM, Seller C, et al. Evaluation of the size 4 i-gel™ airway in one hundred non-paralysed patients*. *Anaesthesia*. 2008;63(10):1124-1130. doi:10.1111/j.1365-2044.2008.05561.x
14. Wharton NM, Gibbison B, Gabbott DA, Haslam GM, Muchatuta N, Cook TM. I-gel insertion by novices in manikins and patients*. *Anaesthesia*. 2008;63(9):991-995. doi:10.1111/j.1365-2044.2008.05542.x
15. Somri M, Matter I, Gaitini LA, Safadi A, Hawash N, Gómez-Ríos MÁ. Fiberoptic-Guided and Blind Tracheal Intubation Through iLTS-D, Ambu® Auragain™, and I-Gel® Supraglottic Airway Devices: A Randomized Crossover Manikin Trial. *J Emerg Med*. 2020;58(1):25-33. doi:10.1016/j.jemermed.2019.09.045
16. Ludena JA, Bellas JJA, Alvarez-Rementeria R, Munoz LE. Fiberoptic-guided intubation after awake insertion of the I-gel™ supraglottic device in a patient with predicted difficult airway. *J Anaesthesiol Clin Pharmacol*. 2017;33(4):560-561. doi:10.4103/0970-9185.173388
17. Levitan RM, Kinkle WC. Initial anatomic investigations of the I-gel airway: a novel supraglottic airway without inflatable cuff*. *Anaesthesia*. 2005;60(10):1022-1026. doi:10.1111/j.1365-2044.2005.04258.x
18. Wang HE, Simeone SJ, Weaver MD, Callaway CW. Interruptions in Cardiopulmonary Resuscitation From Paramedic Endotracheal Intubation. *Ann Emerg Med*. 2009;54(5):645-652.e1. doi:10.1016/j.annemergmed.2009.05.024
19. Lee DW, Kang MJ, Kim YH, et al. Performance of intubation with 4 different airway devices by unskilled rescuers: manikin study. *Am J Emerg Med*. 2015;33(5):691-696. doi:10.1016/j.ajem.2015.03.006
20. Tiesmeier J, Emmerich M. Successful use of the I-gel supraglottic airway as a conduit for blind pre-hospital intubation during resuscitation. *Resuscitation*. 2011;82(10):1358-1359. doi:10.1016/j.resuscitation.2011.07.004

References

21. Theiler L, Kleine-Brueggeney M, Urwyler N, Graf T, Luyet C, Greif R. Randomized clinical trial of the i-gel™ and Magill tracheal tube or single-use ILMA™ and ILMA™ tracheal tube for blind intubation in anaesthetized patients with a predicted difficult airway. *Br J Anaesth*. 2011;107(2):243-250. doi:10.1093/bja/aer102
22. Zamudio-Burbano MA, Giraldo-Salazar O, Gaviria-Rivera E, et al. Tracheal intubation with I-gel supraglottic device in pediatric patients: a prospective case series. *Rev Colomb Anesthesiol*. 2018;46(1):37-41. doi:10.1097/cj9.0000000000000007
23. Anuradha N, Anandhakumar G, David D, Shanmugapriya V. A Prospective Randomized Comparative Study on i-gel versus Laryngeal Mask Airway Fastrach as a Conduit for Blind Tracheal Intubation Using Conventional PVC Endotracheal Tubes in Reverse Orientation. 2017;5(2):6.
24. Crewdson K, Lockey DJ, Røislien J, Lossius HM, Rehn M. The success of pre-hospital tracheal intubation by different pre-hospital providers: a systematic literature review and meta-analysis. *Crit Care*. 2017;21(1):31. doi:10.1186/s13054-017-1603-7
25. Jarvis JL, Wampler D, Wang HE. Association of patient age with first pass success in out-of-hospital advanced airway management. *Resuscitation*. 2019;141:136-143. doi:10.1016/j.resuscitation.2019.06.002